



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the tension in the ring of $P_7N_7O_{14}H_{14}$ is too great to permit of its existence and it spontaneously takes up water, forming $P_7N_7O_{15}H_{16}$. A further confirmation is found in the fact that the higher members on decomposition in part close again to form the stable ring-acid $P_4N_4O_8H_8$, indicating a disposition to form rings containing P_4N_4 . An attempt to test this theory further will be made by endeavoring to prepare diphosphonitric chlorid, $P_2N_2Cl_4$. The corresponding meta-phosphimic acid, $P_2N_2O_4H_4$, should have the angle 90° , differing from the angle of least tension, 135° , by 45° . Such an acid should be much less stable than even $P_7N_7O_{14}H_{14}$ and should pass at once into $P_2N_2O_5H_6$.

WILLIAM H. KRUG,
Secretary.

ACADEMY OF NATURAL SCIENCES OF PHILADELPHIA, MAY 17.

MR. C. S. BOYER illustrated the structure and geographical distribution of diatoms by a large series of lantern views preparatory to describing the following new species: *Rhabdonema Woolmanianum*, *Biddulphia semicircularis Asburyana*, *B. argus*, *B. interrupta*, *B. keeleyi*, *B. Shulzei* and *B. verrucosa*. The paper, with figures, will be published in the *Proceedings* of the Academy.

MR. T. C. PALMER described and illustrated some of the phenomena of conjugation in *Closterium*. The essential steps are essentially as in *Spirogyra*, in that in both cases the process is at first a putting out of tubes which meet and fuse. But just as *Closterium* differs from *Spirogyra* in its method of cell-division, so it presents peculiarities in manner of formation of the zygospore. The two lobes of the desmid seem to possess a certain degree of individuality, at least at the period of conjugation. These lobes, owing to the peculiar method of growth of the plant, generally differ, at the time of conjugation, in age, and therefore in size, and in the thickness, color and markings of the cell-walls. The contrast between two ends or lobes of a given cell is often very great; and in *C. acerosum*, as a rule, each desmid first separates into two entirely distinct and independent semi-cells, each of which is beautifully rounded off at its blunt end by a new growth of cell-wall. The young semi-cell of each desmid then conjugates

with the old semi-cell of the other, and two perfectly distinct zygospores are thus formed. These zygospores and the empty semi-cell cases are held together by a nearly or quite invisible jelly. The 'individuality of the semi-cell,' a tendency toward which has been remarked upon heretofore by Mr. Archer in the case of *C. lineatum*, here becomes practically complete.

In one instance *C. acerosum* formed three zygospores instead of two. One of these was the usual size, made up of the commingled contents of an old and new semi-cell. The other two were about half the size, and consisted each of the unmixed contents of another semi-cell. Of a similar nature is the recently observed discharge, without conjugation, of the contents of a whole cell in *C. lineatum*. The protoplasm, containing small round or ovoid bodies like those in the ordinary zygospore, issued from the ruptured union of the semi-cell cases. It assumed a spherical form. Its development could not be followed further.

The development of the zygospores of *Closterium* is not thoroughly well understood, but the phenomena are probably similar to those of the germination of *Cosmarium*. In addition, however, to this process, another method of reproduction is suspected in *Closterium*, of which the discharge of the cell-contents without conjugation may be one of the stages.

Many of the phases of reproduction in the desmids may be observed to advantage by placing zygospores in life-slides and following the changes that ensue. In such slides large numbers of very minute *Closterium* frequently appear, and these grow perceptibly from day to day, but it is not certain, or even probable, that these arise from the ordinary zygospore.

The following papers were presented for publication in the *Proceedings*:

'Descriptions of five new Phyllostome Bats,' by Gerrit S. Miller, Jr.

'Chitons collected by Dr. Harold Heath at Pacific Grove, near Monterey, Cal.,' by H. A. Pilsbry.

EDWARD J. NOLAN,
Recording Secretary.

Erratum: In the review of Wilder's System of Nomenclature, p. 716, col. 1, line 5, for 'chippocamp' read 'hippocamp.'